

4/6
58. (Once amended) A vector comprising:

- 12
B3
- (a) a first promoter operably linked to an exon defined at its 3' end by an unpaired splice donor site, and
 - (b) a second promoter operably linked to a sequence encoding a selectable marker that lacks an operably linked polyadenylation signal;

wherein said first and second promoters are present in said vector in the same orientation.

4/6
65. (Once amended) A vector comprising a first promoter and a second promoter, said first and second promoters being oriented in the same direction, wherein:

- 12
B3
- (a) said first promoter, but not said second promoter, is operably linked to an exon defined at its 3' end by an unpaired splice donor site; and
 - (b) said vector comprises no operably-linked polyadenylation signals downstream of either said first promoter or said second promoter.

67. (Once amended) A vector comprising:

- B3
- (a) a first promoter operably linked to a sequence encoding a first selectable marker and an unpaired splice donor site; and
 - (b) a second promoter operably linked to a sequence encoding a second selectable marker, wherein neither said first selectable marker

sequence nor said second selectable marker sequence contains an operably-linked polyadenylation signal;

wherein said first and second promoters are present in said vector in the same orientation.

68. (Once amended) The vector of claim 67, wherein said first and second selectable marker sequences are positive selectable marker sequences.

69. (Once amended) The vector of claim 67, wherein said first selectable marker sequence is located upstream of said second selectable marker sequence.

B3
70. (Once amended) A vector construct comprising:

- (COP'4)
- (a) a first promoter operably linked to a sequence encoding a positive selectable marker;
 - (b) a second promoter operably linked to a sequence encoding a negative selectable marker; and
 - (c) an unpaired splice donor site,

wherein said positive and negative selectable marker sequences and said splice donor site are oriented in said vector construct in an orientation such that, when said vector construct is integrated into the genome of a eukaryotic host cell and the vector-encoded splice donor is spliced to a splice acceptor in an endogenous gene in said

B3
C2
genome, then said positive selectable marker sequence is expressed in active form and
said negative selectable marker sequence is not expressed.

B4
73. (Once amended) The vector of any one of claims 58, 65, 67, 70, or 71, said
vector further comprising sequences encoding one or more amplifiable markers.

79. (Once amended) A host cell *in vitro* comprising the vector of any one of
claims 58, 65, 67, 70, or 71.

80. (Once amended) A host cell *in vitro* comprising the vector of claim 72.

B5
81. (Once amended) A host cell *in vitro* comprising the vector of claim 73.

82. (Once amended) A host cell *in vitro* comprising the vector of claim 74.

83. (Once amended) A host cell *in vitro* comprising the vector of claim 75.

84. (Once amended) A host cell *in vitro* comprising the vector of claim 78.

B6
C2
87. (Once amended) A library of cells *in vitro* comprising the vector of any one
of claims 58, 65, 67, 70, or 71.

88. (Once amended) A library of cells *in vitro* comprising the vector of claim 72.

89. (Once amended) A library of cells *in vitro* comprising the vector of claim 73.

90. (Once amended) A library of cells *in vitro* comprising the vector of claim 74.

91. (Once amended) A library of cells *in vitro* comprising the vector of claim 75.

92. (Once amended) A library of cells *in vitro* comprising the vector of claim 78.

93. (Once amended) A method for activation of an endogenous gene in a cell *in vitro* comprising:

(a) transfecting a cell *in vitro* with the vector of any one of claims 58, 65, 67, 70, or 71; and

(b) culturing said cell under conditions suitable for non-homologous integration of said vector into the genome of said cell, wherein said integration results in the activation of an endogenous gene in the genome of said cell.

94. (Once amended) A method for obtaining cDNA from an endogenous gene comprising:

- 100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
- (a) transfecting a plurality of cells *in vitro* with the vector of any one of claims 58, 65, 67, 70, or 71;
 - (b) culturing said cells under conditions suitable for non-homologous integration of the vector into the genome of the cell;
 - (c) selecting for cells in which said vector has integrated into the genomes of said cells;
 - (d) isolating RNA from said selected cells;
 - (e) producing cDNA from said isolated RNA; and
 - (f) isolating one or more cDNA molecules containing one or more nucleotide sequences from said vector.

Sub E1
95. (Once amended) The method of claim 94, wherein said isolation is accomplished by hybridizing said cDNA to said vector.

Sub C1
96. (Once amended) The method of claim 94, wherein said isolation is accomplished by sequencing said cDNA and comparing the nucleotide sequence of said cDNA to the nucleotide sequence of said vector.

97. (Once amended) The vector of claim 67, wherein said unpaired splice donor site is positioned upstream of said first selectable marker sequence such that, when said vector is integrated into the genome of a eukaryotic host cell resulting in splicing from said unpaired splice donor site to a genome-encoded splice acceptor site, then said first selectable marker sequence is not expressed.

98. (Once amended) A method for isolating cells *in vitro* in which a single exon gene has been activated, comprising:

- (a) transfecting a plurality of eukaryotic cells *in vitro* with the vector of claim 97;
- (b) culturing said cells under conditions suitable for non-homologous integration of the vector into the genomes of said cells; and
- (c) selecting for cells in which said first and second selectable marker sequences are expressed in their active forms.

99. (Once amended) A method for isolating a single exon gene cDNA comprising:

- (a) transfecting a plurality of eukaryotic cells *in vitro* with the vector of claim 97;
- (b) culturing said cells under conditions suitable for non-homologous integration of the vector into the genomes of said cells;

- (c) selecting for cells in which said first and second selectable marker sequences are expressed in their active forms;
- (d) isolating RNA from the selected cells;
- (e) producing cDNA from said isolated RNA; and
- (f) isolating a single exon gene from said cDNA.

100. (Once amended) A method for isolating exon I of a gene comprising:

- (a) transfecting one or more eukaryotic cells *in vitro* with the vector of any one of claims 58, 61, 65, or 67;
- (b) culturing said cells under conditions suitable for non-homologous integration of the vector into the genome of said cells;
- (c) selecting for cells in which said vector has transcriptionally activated an endogenous gene containing one or more exons;
- (d) isolating RNA from said selected cells;
- (e) producing cDNA from said isolated RNA;
- (f) recovering a cDNA molecule containing vector sequence and sequence from said endogenous gene; and
- (g) using said endogenous gene sequence to recover exon I of said endogenous gene.

101. (Once amended) A method for expressing a transcript containing exon I of a gene, said method comprising:

- (a) transfecting one or more eukaryotic cells *in vitro* with the vector of any one of claims 58, 61, 65, or 67;
- (b) culturing said cells under conditions suitable for non-homologous integration of the vector into the genome of said cells; and
- (c) culturing said cells under conditions suitable for expression of a transcript containing exon I from an endogenous gene.

B6-sub
(17.10.1) 102. (Once amended) A method for producing a gene product encoded by an endogenous cellular genomic gene, comprising:

- (a) isolating genomic DNA, containing at least one gene, from a eukaryotic cell;
- (b) inserting into or otherwise combining with said isolated genomic DNA, the vector of any one of claims 58, 59, 61, 65, or 67, thereby producing a vector-genomic DNA complex;
- (c) transfecting said vector-genomic DNA complex into a suitable eukaryotic host cell *in vitro*; and
- (d) culturing said host cell under conditions suitable to result in transcription of one or more genes encoded by said vector contained in said vector-genomic DNA complex.

103. (Once amended) A method for isolating a gene sequence comprising:

- (a) isolating genomic DNA, containing at least one gene, from a eukaryotic cell;
- (b) inserting into or otherwise combining with said isolated genomic DNA, the vector of any one of claims 58, 61, 65, or 67, thereby producing a vector-genomic DNA complex;
- (c) transfecting said vector-genomic DNA complex into a suitable eukaryotic host cell *in vitro*;
- (d) culturing said host cell under conditions suitable to result in transcription of one or more genes encoded by said vector contained in said vector-genomic DNA complex;
- (e) isolating RNA produced by said transcription from said host cell;
- (f) producing one or more cDNA molecules from said isolated RNA; and
- (g) recovering one or more cDNA molecules containing vector sequences at the 5' ends of said cDNA molecules, thereby isolating said gene sequence.

106. (Once amended) A method for producing a protein comprising:

- (a) isolating genomic DNA from one or more cells;

- B7
A N-1
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
- (b) inserting into or otherwise combining with said isolated genomic DNA, the vector of any one of claims 58, 61, 65, or 67, thereby producing a vector-genomic DNA complex;
 - (c) transfecting said vector-genomic DNA complex into a suitable host cell *in vitro*; and
 - (d) culturing said cell under conditions suitable to result in protein expression from said genomic DNA contained in said vector-genomic DNA complex.

B7

113. (Once amended) The vector construct of claim 70, wherein said positive selectable marker sequence is selected from the group consisting of a neomycin gene, a hypoxanthine phosphoribosyl transferase gene, a puromycin gene, a dihydroorotase gene, a glutamine synthetase gene, a histidine D gene, a carbamyl phosphate synthase gene, a dihydrofolate reductase gene, a multidrug resistance I gene, an aspartate transcarbamylase gene, a xanthine-guanine phosphoribosyl transferase gene, and an adenosine deaminase gene.

114. (Once amended) The vector construct of claim 70, wherein said negative selectable marker sequence is selected from the group consisting of a hypoxanthine phosphoribosyl transferase gene, a thymidine kinase gene, and a diphtheria toxin gene.

115. (Once amended) The vector of claim 70, wherein said negative selectable marker sequence is located upstream of said positive selectable marker.

116. (Once amended) The vector of claim 115, wherein said vector further comprises one or more selectable marker sequences.

Please add the following new claims:

117. (New) The vector of claim 67, wherein said unpaired splice donor site is positioned within said first selectable marker sequence such that, when said vector is integrated into the genome of a eukaryotic host cell, resulting in splicing from said unpaired splice donor site to a genome-encoded splice acceptor site, then said first selectable marker sequence is expressed in inactive form.

118. (New) A vector construct comprising:

- (a) a first promoter operably linked to a sequence encoding a positive selectable marker;
- (b) a second promoter operably linked to a sequence encoding a negative selectable marker; and
- (c) an unpaired splice donor site,

wherein said positive and negative selectable marker sequences and said splice donor site are oriented in said vector construct in an orientation such that, when said vector construct is integrated into the genome of a eukaryotic host cell and the vector-encoded splice donor is spliced to a splice acceptor in an endogenous gene in said genome, then said positive selectable marker sequence is expressed in active form and said negative selectable marker sequence is expressed in inactive form.